

# GASTROENTEROLOGY ASSOCIATES OF CENTRAL JERSEY, P.A.

1931 OAK TREE ROAD, SUITE 202 EDISON, N.J. 08820

PHONE 732-744-9090 FAX 732-744-1592 Website: [www.gacj.net](http://www.gacj.net)

ANDREW S. BAIK, M.D. DANIEL A. BAIK, M.D. MEET PARIKH, D.O.

## PLENVU SPLIT DOSE PREP COLONOSCOPY PREPARATION INSTRUCTIONS

---

DATE: \_\_\_\_\_ REGISTRATION TIME: \_\_\_\_\_

**LOCATION:** Oak Tree Surg. Ctr., 1931 Oak Tree Rd. Edison, N.J. 08820 Main Level  
Please call the surgery center upon arrival in the parking lot for further instructions.  
(732)603-8603  
Saint Peter's University Hospital 254 Easton Ave, New Brunswick, NJ 08901

The following items *must be purchased from the pharmacy* for your preparation:

- ONE (1)** PLENVU KIT for Oral Solution [with prescription] – this will be mailed from pharmacy  
**TWO (2)** EXTRA STRENGTH SIMETHICONE tablets 125mg (i.e. Gas-X or generic) [over-the-counter]

### SEVEN (7) DAYS BEFORE EXAMINATION:

- Stop all **WEIGHT LOSS MEDICATIONS** (i.e. Ozempic, Wegovy, Mounjaro, Saxenda). It is ok to take these medications following completion of your procedure(s).

### FIVE (5) DAYS BEFORE EXAMINATION:

- Stop all **NSAID MEDICATIONS** (i.e. Aspirin, Aleve, Advil, Motrin, Bayer, Ecotrin, Ibuprofen, Naproxen, etc.). It is OK to continue baby aspirin if you take this medication on a daily basis.
- If you take **BLOOD THINNERS** (i.e. Coumadin, Plavix, Eliquis, Xarelto, Heparin, etc.), please review instructions discussed at your office visit and with your primary care physician or cardiologist. Usually, medications like Coumadin and Plavix are held five days prior to the procedure. Medications like Eliquis and Xarelto are usually held two-four days prior to the procedure.
- **TYLENOL** (and other brands containing ACETAMINOPHEN) is **OK** as a pain reliever.

**\*\* ALL OF THE ABOVE SHOULD BE FOLLOWED TO MINIMIZE RISK OF  
SERIOUS BLEEDING IF A POLYP IS REMOVED \*\***

### **THREE (3) DAYS BEFORE EXAMINATION:**

Eat a **LOW RESIDUE DIET**. A Low residue diet limits high fiber food.

**HIGH FIBER FOODS TO AVOID INCLUDE:**

- Whole grain breads, oatmeal/cereals, granola
- Nuts, seeds, raw/dried vegetables or fruit (NO salads)
- Beverages with pulp
- Nutritional supplements that contain fiber
- Peppers, beans, corn/popcorn

### **DAY BEFORE & DAY OF EXAMINATION:**

We encourage you to stay hydrated with at least 12 tall glasses (about 8-10 ounces each) of clear liquids throughout the day before your procedure in addition to what you drink with your bowel prep medication to prevent dehydration.

Stay on a **CLEAR LIQUID DIET** throughout the entire day before your procedure:

**\*NOTHING RED OR PURPLE\***

Things that are OK to drink/eat:

- Strained **clear** fruit juice **without pulp**
- Water, Gatorade, PowerAde, soft drinks (ginger ale, sprite)
- Coffee or tea **without milk or creamer**
- Plain Jell-o **without fruit or toppings**
- Ice popsicles (no sherbet or fruit bars)
- Fat free / Low sodium clear broth (nothing greasy or chunks)

**\*\*Do NOT eat any solid food while on this diet. Do NOT drink alcohol. NO dairy products.\*\***

At **6:00 p.m. the night before your procedure** using the mixing container, add **Dose 1** with at least 16 ounces of water. Shake vigorously until it is completely dissolved. Drink the solution slowly, optimally within 30 minutes to 1 hour. Then, using your container, drink at least an additional 16 ounces of water and try to finish this within an hour. Continue to drink additional clear liquids until your second dose.

**7 HOURS** prior to your procedure arrival time at \_\_\_\_\_, take 2 Extra Strength Simethicone tablets (total of 250mg) and then begin Dose 2 of the prep. Add **Dose 2 (both Pouch A and Pouch B)** in the container with at least 16 ounces of water. Shake vigorously until it is completely dissolved. Drink your solution slowly, optimally within 30 minutes to 1 hour. Then, using your container, drink at least an additional 16 ounces of water within an hour.

**STOP DRINKING ALL CLEAR LIQUIDS BY \_\_\_\_\_ (5 HOURS BEFORE** your procedure arrival time). This means that you should not have anything to eat or drink 5 hours and onwards before you arrive for your procedure. We encourage you to drink plenty of fluids until this time. If you take any blood pressure or heart medication, please take your regular dose with a very small sip of water.

- You will develop significant diarrhea after drinking the preparation. Plan to be near a bathroom. This is normal as it means the medication is working to clear stool from your colon.

- Most people feel mild bloating and mild abdominal cramps. This is normal. Drinking the prep medication more slowly and over a longer period can help alleviate these symptoms.
- A successful colon prep will cause you to have clear yellow (“tea-colored”) liquid stools. Please finish all your preparation regardless of your stool color.

- **NO NSAID MEDICATIONS** (i.e. Aspirin, Aleve, Advil, Motrin, Bayer, Ecotrin, Ibuprofen, Naproxen, etc.) **FIVE DAYS PRIOR TO TEST.**  
It is OK to continue baby aspirin if you take this medication on a daily basis.
- **IF YOU TAKE BLOOD THINNERS**, please ensure you followed all instructions for stopping them prior to the procedure.
- **TYLENOL** (and other brands which contain ACETAMINOPHEN) IS **O.K. TO TAKE!!!**
- **YOU MUST HAVE SOMEONE TO DRIVE YOU HOME.**
- **PLEASE LEAVE ALL THE JEWELRY AT HOME.**
- **PLEASE BRING YOUR INSURANCE CARD & PHOTO ID.**
- **PLEASE BRING A LIST OF YOUR MEDICATIONS WITH DOSAGES.**
- **PLEASE BRING THE COMPLETED PATIENT INFORMATION PACKET (IF NEEDED).**

#### Commonly asked Questions:

- ***How do I know if my prep is adequate?***  
The stool should be watery in consistency. It does not have to be clear in color like water since digestive juices will continue to tint the stool yellow and small flecks of debris are not a problem as long as the stool is not muddy or thick.
- ***What side effects may I expect?***  
Since the prep works by flooding the intestinal tract with fluid, abdominal bloating and cramping may occur, as well as some nausea and vomiting. This is usually temporary, and as the diarrhea develops, symptoms will gradually improve. Weaknesses and lightheadedness can also occur, especially if you have not taken enough fluid with electrolytes with the prep, so be sure to have drinks such as Gatorade/Powerade and Pedialyte.
- ***Will the prep interfere with my other medications?***  
Medications taken at least an hour before beginning the prep should be adequately absorbed, but thereafter, they are likely to be washed away by the prep.
- ***What if I have other questions?***  
Call the office at 732-744-9090

Please be aware that this procedure will be taking place at an ambulatory surgical center. Therefore, please be advised that all of your AMBULATORY SURGICAL IN-NETWORK BENEFITS such as co-pays, deductibles, and co-insurances will apply. We strongly recommend that you contact your insurance company to most accurately verify your benefits.

**This is NOT an office procedure.**