

GASTROENTEROLOGY ASSOCIATES OF CENTRAL JERSEY, P.A.

1931 OAK TREE ROAD, SUITE 202 EDISON, N.J. 08820

PHONE 732-744-9090 FAX 732-744-1592 Website: www.gacj.net

ANDREW S. BAIK, M.D. DANIEL A. BAIK, M.D. MEET PARIKH, D.O.

GOLYTELY

COLONOSCOPY PREPARATION INSTRUCTIONS

DATE: _____ REGISTRATION TIME: _____

LOCATION: Oak Tree Surg. Ctr., 1931 Oak Tree Rd. Edison, N.J. 08820 Main Level
Please call the surgery center upon arrival in the parking lot for further instructions.
(732)603-8603

Saint Peter's University Hospital 254 Easton Ave, New Brunswick, NJ 08901

The following items ***must be purchased*** your local pharmacy for your preparation:

- **ONE (1)** GOLYTELY Bowel Prep Kit [with prescription]
- **TWO (4)** BISACODYL tablets 5mg (i.e. Dulcolax or generic brand) [over-the-counter]

SEVEN (7) DAYS BEFORE EXAMINATION:

- Stop all **WEIGHT LOSS MEDICATIONS** (i.e. Ozempic, Wegovy, Mounjaro, Saxenda). It is ok to take these medications following completion of your procedure(s).

FIVE (5) DAYS BEFORE EXAMINATION:

- Stop all **NSAID MEDICATIONS** (i.e. Aspirin, Aleve, Advil, Motrin, Bayer, Ecotrin, Ibuprofen, Naproxen, etc.). It is OK to continue baby aspirin if you take this medication on a daily basis.
- If you take **BLOOD THINNERS** (i.e. Coumadin, Plavix, Eliquis, Xarelto, Heparin, etc.), please review instructions discussed at your office visit and with your primary care physician or cardiologist. Usually, medications like Coumadin and Plavix are held five days prior to the procedure. Medications like Eliquis and Xarelto are usually held two-four days prior to the procedure.
- **TYLENOL** and other brands containing **ACETAMINOPHEN** is **OK** as a pain reliever.

**** ALL OF THE ABOVE SHOULD BE FOLLOWED TO MINIMIZE RISK OF SERIOUS BLEEDING IF A POLYP IS REMOVED ****

THREE (3) DAYS BEFORE EXAMINATION:

- Eat a **LOW RESIDUE DIET**. A Low residue diet limits high fiber food.
- **HIGH FIBER FOODS TO AVOID INCLUDE:**
 - Whole grain breads, oatmeal/cereals, granola
 - Nuts, seeds, raw/dried vegetables or fruit (NO salads)
 - Beverages with pulp
 - Nutritional supplements that contain fiber
 - Peppers, beans, corn/popcorn

DAY BEFORE EXAMINATION:

We encourage you to stay hydrated with at least 12 tall glasses (about 8-10 ounces each) of clear liquids throughout the day in addition to what you drink with your bowel prep medication to prevent dehydration.

Have a *clear* LIQUID diet throughout the entire day: ** **NOTHING RED OR PURPLE** **

Things that are OK to drink/eat:

- Strained **clear** fruit juice **without pulp**
- Water, Gatorade, PowerAde, soft drinks (ginger ale, sprite)
- Coffee or tea **without milk or creamer**
- Plain jell-o **without fruit or toppings**
- Ice popsicles (no sherbet or fruit bars)
- Fat free / Low sodium clear broth (nothing greasy or chunks)

Do **NOT** eat any solid food while on this diet. Do **NOT** drink alcohol. **NO** dairy products.

1. At **12:00 p.m.**, take 4 Bisacodyl tablets with a glass of. **DO NOT** chew or crush.
2. Between **3-6 p.m.** the night before your procedure, shake vigorously again and drink half of the bottle (2 liters): one, **10 oz. glass of GoLyteLy solution every 10-15 minutes**. If you are able, walk about while drinking the liquid.
3. At **10:00 p.m.**, drink the second half of the bottle (2 liters); one, 10oz. **glass of GoLyteLy solution** every 10-15 minutes. Drink each glass quickly rather than drinking small amounts continuously, using a drinking straw may help to drink the liquid quickly.
Continue drinking clear liquids until midnight.

DAY OF EXAMINATION:

STOP DRINKING ALL CLEAR LIQUIDS 8 HOURS BEFORE your registration time. This means that you should not have anything to eat or drink 8 hours before your colonoscopy and onward. We encourage you to drink plenty of fluids until this time. If you take any blood pressure or heart medication, please take your regular dose with very small sips of water.

- ❖ **NO NSAID MEDICATIONS** (i.e. Aspirin, Aleve, Advil, Motrin, Bayer, Ecotrin, Ibuprofen, Naproxen, etc.) **FIVE DAYS PRIOR TO TEST.**
It is OK to continue baby aspirin if you take this medication on a daily basis.
- ❖ **IF YOU TAKE BLOOD THINNERS**, please ensure you followed all instructions for stopping them prior to the procedure.
- ❖ **TYLENOL** (and other brands which contain ACETAMINOPHEN) IS **O.K. TO TAKE!!!**
- ❖ YOU **MUST** HAVE SOMEONE TO **DRIVE YOU HOME.**
- ❖ PLEASE LEAVE ALL **JEWELRY AT HOME.**
- ❖ PLEASE BRING YOUR **INSURANCE CARD & PHOTO ID.**
- ❖ PLEASE BRING **THE COMPLETED PATIENT INFORMATION PACKET (IF NEEDED).**

Commonly asked Questions:

- ***How do I know if my prep is adequate?***
The stool should be watery in consistency. It does not have to be clear in color like water since digestive juices will continue to tint the stool yellow and small flecks of debris are not a problem as long as the stool is not muddy or thick.
- ***What side effects may I expect?***
Since the prep works by flooding the intestinal tract with fluid, abdominal bloating and cramping may occur, as well as some nausea and vomiting. This is usually temporary, and as the diarrhea develops, symptoms will gradually improve. Weaknesses can also occur, especially if you have not taken enough fluid with the prep, and can be remedied by increasing fluid intake.
- ***Will the prep interfere with my other medications?***
Medications taken at least an hour before beginning the prep should be adequately absorbed, but thereafter, they are likely to be washed away by the prep.
- ***What if I have other questions?***
Call the office at 732-744-9090

Please be aware that this procedure will be taking place at an ambulatory surgical center. Therefore, please be advised that all of your **AMBULATORY SURGICAL IN-NETWORK BENEFITS** such as co-pays, deductibles, and co-insurances will apply. We strongly recommend that you contact your insurance company to most accurately verify your benefits.

This is NOT an office procedure.